arXiv:1408.1074v1 [math.HO] 5 Aug 2014

Appell F_1 and Conformal Mapping

Steven Finch

August 5, 2014

Abstract. This is the last of a trilogy of papers on triangle centers. A fairly obscure "conformal center of gravity" is computed for the class of all isosceles triangles. This calculation appears to be new. A byproduct is the logarithmic capacity or transfinite diameter of such, yielding results consistent with Haegi (1951).

Before discussing triangles, let us give both a review of [1] and a preview involving a simpler region in the plane.

Let $\Omega = \{x + iy \in \mathbb{C} : y > 0,\ x^2 + y^2 < 1\}$, the interior of the upper half-disk of unit radius. Let Δ denote the (full) disk of unit radius and Σ denote the infinite horizontal strip of width π. Define a function $\ell : \Omega \to \Sigma$ by [2]

$$\ell(z) = \ln\left(\frac{(1+z)^2}{(1-z)^2}\right).$$

The conformal map $f_w : \Omega \to \Delta$ given by

$$f_w(z) = \frac{\exp(\ell(z)) - \exp(\ell(w))}{\exp(\ell(z)) - \exp(\overline{\ell(w)})}$$

satisfies $f_w(w) = 0$; it is well-known that $\ln|f_w(z)|$ is Green's function for Ω. We deduce that

$$h(w) = \lim_{z \to w}\left|\frac{f_w(z)}{z-w}\right| = \left|\frac{\exp(\ell(w))\ell'(w)}{\exp(\ell(w)) - \exp(\overline{\ell(w)})}\right|$$

where ℓ' denotes the derivative of ℓ. Restricting attention to the y-axis only, we have

$$h(iy) = \left|\frac{\frac{2(1+iy)}{(1-iy)^2} + \frac{2(1+iy)^2}{(1-iy)^3}}{\frac{(1+iy)^2}{(1-iy)^2} - \frac{(i+y)^2}{(-i+y)^2}}\right| = \left|\frac{(1+iy)^2}{2y\,(1-y^2)}\right| = \frac{1+y^2}{2y\,(1-y^2)}$$

for $0 < y < 1$. Minimizing this expression, it follows that

$$iy_0 = i\sqrt{-2+\sqrt{5}} = (0.485868271756645678182863 8...)i$$

[0]Copyright © 2014 by Steven R. Finch. All rights reserved.

is the **inner conformal center** (what was called the least capacity point in [1]) for Ω. Also,

$$\frac{1}{h(iy_0)} = \sqrt{-22 + 10\sqrt{5}} = 0.60056621200155521577338 94...$$

is the **maximum inner radius** of Ω [3, 4]. This concludes our review.

Let Ω^c denote the complement of the closure of Ω. Inverting a function [5]

$$\frac{1}{z} = \frac{\left(\frac{\sqrt{3}}{2} - \frac{i}{2}\right)\left(1 + \frac{1}{w}\right)^{2/3} + \left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right)\left(1 - \frac{1}{w}\right)^{2/3}}{\left(1 + \frac{1}{w}\right)^{2/3} - \left(1 - \frac{1}{w}\right)^{2/3}}$$

in terms of w yields

$$\frac{\left(1 + \frac{1}{w}\right)^{2/3}}{\left(1 - \frac{1}{w}\right)^{2/3}} = \frac{\frac{\sqrt{3}}{2} + \frac{i}{2} + \frac{1}{z}}{-\frac{\sqrt{3}}{2} + \frac{i}{2} + \frac{1}{z}} = m(z).$$

The conformal map $g : \Delta \to \Omega^c$

$$g(z) = \frac{1 + 2m(z)^{3/2} + m(z)^3}{-1 + m(z)^3} = \frac{4}{3\sqrt{3}}\frac{1}{z} + \frac{2i}{3\sqrt{3}} + O(z)$$

satisfies $g(0) = \infty$, has positive leading Laurent coefficient, and is unique in this regard. The constant term of the series expansion

$$\frac{2i}{3\sqrt{3}} = (0.38490017945975050967276 58...)i$$

is the **outer conformal center**, which clearly lies in Ω but is not the same as the inner conformal center. A motivating feature is

$$\frac{1}{2\pi}\int_0^{2\pi} g\left((1-\varepsilon)e^{it}\right) dt = \frac{2i}{3\sqrt{3}}$$

for $\varepsilon > 0$, but the literature is small [6, 7, 8, 9, 10]. The leading coefficient (of $1/z$) is the **outer radius** of Ω [3, 11]:

$$\frac{4}{3\sqrt{3}} = 0.76980035891950101934553 17...$$

which is also known as the **logarithmic capacity** or **transfinite diameter** of Ω. An alternative definition is [12]

$$\lim_{n\to\infty} \max_{z_1,z_2,\ldots,z_n\in\Omega} \left(\prod_{j<k}|z_j - z_k|\right)^{\frac{2}{n(n-1)}},$$

that is, the maximal geometric mean of pairwise distances for n points in Ω, in the limit as $n \to \infty$. This constitutes a fascinating collision of ideas from potential theory; replacing a geometric mean by an arithmetic mean seems to be an open computational issue.

As a conclusion to our preview, the Appell F_1 function can be written as a definite integral [13, 14]

$$F_1(a,b,b',c;x,y) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)}\int_0^1 s^{a-1}(1-s)^{c-a-1}(1-sx)^{-b}(1-sy)^{-b'}ds,$$

$$|x|<1, \qquad |y|<1, \qquad \mathrm{Re}(c)>\mathrm{Re}(a)>0$$

as well as a double hypergeometric series

$$F_1(a,b,b',c;x,y) = \sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\frac{1}{m!n!}\frac{\Gamma(a+m+n)}{\Gamma(a)}\frac{\Gamma(b+m)}{\Gamma(b)}\frac{\Gamma(b'+n)}{\Gamma(b')}\frac{\Gamma(c)}{\Gamma(c+m+n)}x^m y^n.$$

Its numerical implementation in Mathematica is crucial to everything that follows.

1. Isosceles Triangles

Let

$$T_\theta = \{x+iy \in \mathbb{C} : 0<x<\cos(\theta/2),\ \ y<\tan(\theta/2)x,\ \ y>-\tan(\theta/2)x\},$$

the interior of an isosceles triangle with apex angle $0<\theta<\pi$ located at the origin. Two sides of unit length meet there; the third (vertical) side has x-intercept $\cos(\theta/2)$ and length $2\sin(\theta/2)$.

To construct a conformal map $g:\Delta\to T_\theta^c$ requires two steps. First, define

$$f(z) = \int_{z_0}^{z}\frac{(\zeta-a_1)^{\mu_1}(\zeta-a_2)^{\mu_2}(\zeta-a_3)^{\mu_3}}{\zeta^2}d\zeta$$

on Δ, where a_1, a_2, a_3 are *prevertices* of the unit circle mapping onto vertices of T_θ, and $\pi(1+\mu_1)$, $\pi(1+\mu_2)$, $\pi(1+\mu_3)$ are exterior angles at the corresponding vertices.

Also, z_0 is some point of Δ other than 0, and the integral is taken along any curve in Δ joining z_0 to z not passing through 0 (it does not matter which). Clearly

$$\mu_1 = \frac{\pi - \theta}{\pi}, \qquad \mu_2 = \mu_3 = \frac{\pi + \theta}{2\pi}.$$

The choice of point a_1 is arbitrary; here let $a_1 = -1$. The remaining two points a_2, a_3 must satisfy the constraint [15, 16]

$$\mu_1/a_1 + \mu_2/a_2 + \mu_3/a_3 = 0$$

in order that $f(0) = \infty$. Thus

$$a_2 = \frac{\pi - \theta + 2i\sqrt{\pi\theta}}{\pi + \theta}, \qquad a_3 = \frac{\pi - \theta - 2i\sqrt{\pi\theta}}{\pi + \theta}$$

work for our purposes. We further choose $z_0 = -1$, so that

$$f(z) = \int_{-1}^{z} \frac{(\zeta+1)^{1-\theta/\pi}(\zeta^2 - 2\,\frac{\pi-\theta}{\pi+\theta}\,\zeta + 1)^{(\pi+\theta)/(2\pi)}}{\zeta^2} d\zeta.$$

The image of $\{a_1, a_2, a_3\}$ under f evidently lies in the left half plane – needing rotation by π – plus rescaling so that the vertical triangle side has the proper length. This second step is achieved by defining

$$g(z) = -\frac{2\sin(\theta/2)}{\operatorname{Im} f(a_2) - \operatorname{Im} f(a_3)} f(z).$$

For the scenario $\theta = \pi/2$, it is true that the coefficient

$$-\frac{\sqrt{2}}{\operatorname{Im} f(a_2) - \operatorname{Im} f(a_3)} = 0.47563444387998193205675 70... = \frac{3^{3/4}}{2^{7/2}\pi^{3/2}}\Gamma\left(\frac{1}{4}\right)^2 = \kappa$$

to high numerical precision. This expression (the outer radius of an isosceles right triangle) is well-known and is a special case of a more general formula due to Haegi [3, 17]. More on this will be given soon.

Our key result is that the function $f(z)$ possesses an exact representation. Let

$$\xi(z) = \frac{\left(\sqrt{\pi} - i\sqrt{\theta}\right)(z+1)}{2\sqrt{\pi}}, \qquad \eta(z) = \frac{\left(\sqrt{\pi} + i\sqrt{\theta}\right)(z+1)}{2\sqrt{\pi}},$$

$$\varphi(z) = \left[\frac{\pi\,(z-1)^2 + \theta(z+1)^2}{\pi+\theta}\right]^{(\pi+\theta)/(2\pi)},$$

$$\psi(z) = \left[\frac{\pi\,(z-1)^2 + \theta(z+1)^2}{\pi+\theta}\right]^{(\pi-\theta)/(2\pi)},$$

$$\delta(z) = \left[-\sqrt{\pi}\,(z-1) - i\sqrt{\theta}(z+1)\right]^{(\pi-\theta)/(2\pi)} \left[-\sqrt{\pi}\,(z-1) + i\sqrt{\theta}(z+1)\right]^{(\pi-\theta)/(2\pi)}.$$

Then we have

$$f(z) = (z+1)^{1-\theta/\pi}\left\{-\frac{\varphi(z)}{z} + \frac{2^{\theta/\pi}\pi^{(\pi+\theta)/(2\pi)}}{2\pi^2+\pi\theta-\theta^2}\frac{\delta(z)}{\psi(z)}\cdot \right.$$
$$\left[-2(2\pi-\theta)F_1\left(1-\frac{\theta}{\pi}, \frac{\pi-\theta}{2\pi}, \frac{\pi-\theta}{2\pi}, 2-\frac{\theta}{\pi}; \xi(z), \eta(z)\right) + \right.$$
$$\left.\left.(\pi+\theta)(z+1)F_1\left(2-\frac{\theta}{\pi}, \frac{\pi-\theta}{2\pi}, \frac{\pi-\theta}{2\pi}, 3-\frac{\theta}{\pi}; \eta(z), \xi(z)\right)\right]\right\}$$

as can be easily proved after-the-fact by differentiation. (Our before-the-fact technique consisted of examining rational multiples of θ in Mathematica, seeking recognizable patterns.) This integral evaluation appears to be new.

Returning to the $\theta = \pi/2$ scenario,

$$g(z) = \frac{\kappa}{z} + \lambda + O(z)$$

as $z \to 0$, where

$$\lambda = \kappa\frac{2^{5/4}}{3^{3/4}}\left[2F_1\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{2}; \frac{2-i\sqrt{2}}{4}, \frac{2+i\sqrt{2}}{4}\right) - F_1\left(\frac{3}{2}, \frac{1}{4}, \frac{1}{4}, \frac{5}{2}; \frac{2+i\sqrt{2}}{4}, \frac{2-i\sqrt{2}}{4}\right)\right]$$
$$= 0.5045039334500261012764068...$$

is the outer conformal center of $T_{\pi/2}$. We wonder whether this expression for λ can be simplified, for example, as a ratio of gamma or Gauss hypergeometric function values.

As a corollary, let $\tilde{T} = \{x+iy \in \mathbb{C} : x > 0,\, y > 0,\, x+y < 1\}$, the initial triangle examined in [1, 18]. The outer conformal center of $\tilde{T}$ is simply

$$\left(\frac{1+i}{\sqrt{2}}\right)\lambda = (0.35673815247780014067513 07...)(1+i)$$

which is not the same as the inner conformal center $(0.301...)(1+i)$.

We mention finally that the outer conformal center of $T_{\pi/3}$ (an equilateral triangle) is $1/\sqrt{3}$, that is, it coincides with the centroid of $T_{\pi/3}$. No other scenarios with such recognizable λ have been found.

2. HAEGI'S FORMULA

An arbitrary triangle with sides a, b, c and opposite angles

$$\alpha = \arccos\left(\frac{b^2+c^2-a^2}{2bc}\right),$$

$$\beta = \arccos\left(\frac{a^2+c^2-b^2}{2ac}\right),$$

$$\gamma = \arccos\left(\frac{a^2+b^2-c^2}{2ab}\right)$$

has area, circumradius and logarithmic capacity given by

$$A = \sqrt{\frac{a+b+c}{2}\frac{-a+b+c}{2}\frac{a-b+c}{2}\frac{a+b-c}{2}},$$

$$R = \frac{abc}{\sqrt{(a+b+c)(b+c-a)(c+a-b)(a+b-c)}},$$

$$\kappa = \frac{A}{4\pi^2 q(\alpha/\pi)q(\beta/\pi)q(\gamma/\pi)R}$$

where

$$q(x) = \frac{1}{\Gamma(x)}\sqrt{\frac{x^x}{(1-x)^{1-x}}}.$$

Under the special circumstances that $a = b = 1$ and $c = 2\,\sin(\theta/2)$, we have

$$\kappa(\theta) = \frac{\sqrt{\pi+\theta}}{8\pi^{5/2}}\left(\frac{\pi+\theta}{4\theta}\right)^{\theta/(2\pi)}\frac{\sin(\theta)^2}{\sin(\theta/2)}\Gamma\left(\frac{\theta}{\pi}\right)\Gamma\left(\frac{\pi-\theta}{2\pi}\right)^2$$

for the isosceles triangles T_θ. Over such triangles, the one with maximal κ has $\theta = 2.5360873621...$, which seems not to have been noticed before. Over the family of *all* triangles with fixed A, the one with minimal κ is equilateral, as proved by Pólya & Szegö [3, 19, 20]. If we fix perimeter rather than area, then (to the contrary) the equilateral triangle provides *maximal* κ.

3. ADDENDUM: 30°-60°-90° TRIANGLE

Define $T = \left\{x+iy \in \mathbb{C} : x > 0,\, y > 0,\, \sqrt{3}x + y < \sqrt{3}\right\}$. Proceeding as before, we obtain

$$\mu_1 = \frac{1}{2}, \qquad \mu_2 = \frac{2}{3}, \qquad \mu_3 = \frac{5}{6}.$$

The choice of point a_1 is arbitrary; here let $a_1 = 1$. From $\mu_1/a_1 + \mu_2/a_2 + \mu_3/a_3 = 0$, we deduce that

$$a_2 = i, \qquad a_3 = -\frac{3}{5} - \frac{4}{5}i$$

work for our purposes. Choosing $z_0 = 1$, it follows that

$$f(z) = \int_1^z \frac{(\zeta-1)^{1/2}(\zeta-i)^{2/3}\left(\zeta + \left(\frac{3}{5} + \frac{4}{5}i\right)\right)^{5/6}}{\zeta^2} d\zeta.$$

An exact representation for $f(z)$ in terms of the Appell F_1 function is possible. From this, we can verify the outer radius expression [3, 17]

$$\kappa = \frac{5^{5/12}}{2^{10/3}\pi^2}\Gamma\left(\frac{1}{3}\right)^3 = 0.37791374297095583210248 82...$$

to high numerical precision, but have not yet determined the outer conformal center of T.

4. ADDENDUM: 6-9-13 TRIANGLE

The Schwarz-Christoffel toolbox for Matlab [21, 22] makes numerical computations of a conformal map feasible. For the triangle with vertices

$$0, \quad 6, \quad -\frac{13}{3} + \frac{4\sqrt{35}}{3}i$$

the following code:

```
p = polygon([0 6 -13/3+(4*sqrt(35)/3)*i])
f = extermap(p,scmapopt('Tolerance',1e-18))
p = parameters(f)
format long
p.prevertex
```

gives

$$\mu_1 = 0.659, \qquad \mu_2 = 0.207, \qquad \mu_3 = 0.132,$$

$$a_1 = 1, \qquad a_2 = 0.0163 - 0.9998i, \qquad a_3 = -0.4069 + 0.9134i.$$

Closed-form expressions for these exponents and prevertices are possible yet cumbersome. The same is true for the outer radius $\kappa = 3.805336$. Determining the outer conformal center (even approximately) remains open. Figures 1, 2, 3 provide conformal map images of ten evenly-spaced concentric circles in the disk; orthogonal trajectories are also indicated. We leave the task of exploring whether outer conformal centers belong in Kimberling's database [23] to someone else.



Figure 1: Images of ten concentric circles, center at $0.356 + (0.356)i$.



Figure 2: Images of ten concentric circles, center unknown.



Figure 3: Images of ten concentric circles, center unknown.

5. Acknowledgements

I am grateful to Thomas Ransford [12, 16] for explaining Schwarz-Christoffel prevertex calculations for the scenario $\theta = \pi/2$.

References

[1] S. Finch, Least capacity point of triangles, http://arxiv.org/abs/1407.4105.

[2] J. Kevorkian, *Partial Differential Equations. Analytical Solution Techniques*, 2nd ed., Springer-Verlag, 2000, pp. 129–130; MR1728947 (2000i:35001).

[3] G. Pólya and G. Szegö, *Isoperimetric Inequalities in Mathematical Physics*, Princeton Univ. Press, 1951, pp. 1–3, 158, 254–258, 273–274; MR0043486 (13,270d).

[4] M. Hantke, *Summen reziproker Eigenwerte*, Ph.D. thesis, Martin-Luther-Universität Halle-Wittenberg, 2006, http://sundoc.bibliothek.uni-halle.de/diss-online/06/06H308/index.htm.

[5] Ch. Pommerenke, On metric properties of complex polynomials, *Michigan Math. J.* 8 (1961) 97–115; MR0151580 (27 #1564).

[6] G. Pólya and G. Szegő, *Problems and Theorems in Analysis. I, Series, Integral Calculus, Theory of Functions*, Springer-Verlag, 1998, pp. 129–130, 325–326; MR1492447.

[7] G. Pólya and G. Szegő, *Problems and Theorems in Analysis. II, Theory of Functions, Zeros, Polynomials, Determinants, Number Theory, Geometry*, Springer-Verlag, 1998, pp. 23–24, 192–194; MR1492448.

[8] Ch. Pommerenke, *Univalent Functions*, Vandenhoeck & Ruprecht, 1975, pp. 12–13; MR0507768 (58 #22526).

[9] R. Bouffanais, G. D. Weymouth and D. K. P. Yue, Hydrodynamic object recognition using pressure sensing, *Proc. Royal Soc. London Ser. A* 467 (2011) 19–38; MR2764670.

[10] G. Cleanthous, Monotonicity theorems for analytic functions centered at infinity, *Proc. Amer. Math. Soc.* 142 (2014) 3545–3551.

[11] S. Liesipohja, *Numerical Methods for Computing Logarithmic Capacity*, M.Sc. thesis, University of Helsinki, 2014, https://helda.helsinki.fi/handle/10138/44698.

[12] T. Ransford, *Potential Theory in the Complex Plane*, Cambridge Univ. Press, 1995, pp. 132–137, 152–160; MR1334766 (96e:31001).

[13] W. N. Bailey, *Generalized Hypergeometric Series*, Cambridge Univ. Press, 1935, pp. 73–83.

[14] J. McDougall, L. Schaubroeck and J. Rolf, *Exploring Complex Analysis*. Ch. 5: *Mappings to Polygonal Domains*, http://www.jimrolf.com/explorationsInComplexVariables.html.

[15] Z. Nehari, *Conformal Mapping*, Dover, 1975, pp. 189–195; MR0377031 (51 #13206).

[16] T. Ransford, Logarithmic capacity for $\tilde{T}$, unpublished note (2014).

[17] H. R. Haegi, Extremalprobleme und Ungleichungen konformer Gebietsgrössen, *Compositio Math.* 8 (1950) 81–111; MR0039811 (12,602b).

[18] S. Finch, In limbo: Three triangle centers, http://arxiv.org/abs/1406.0836.

[19] G. Pólya and G. Szegő, Inequalities for the capacity of a condenser, *Amer. J. Math.* 67 (1945) 1–32; MR0011871 (6,227e).

[20] A. Yu. Solynin and V. A. Zalgaller, An isoperimetric inequality for logarithmic capacity of polygons, *Annals of Math.* 159 (2004) 277–303; MR2052355 (2005a:31002).

[21] T. A. Driscoll and L. N. Trefethen, *Schwarz-Christoffel Mapping*, Cambridge Univ. Press, 2002, pp. 9–18; MR1908657 (2003e:30012).

[22] T. A. Driscoll, Schwarz-Christoffel Toolbox for Matlab, http://www.math.udel.edu/~driscoll/SC/.

[23] C. Kimberling, Encyclopedia of Triangle Centers, http://faculty.evansville.edu/ck6/encyclopedia/.

Steven Finch
Dept. of Statistics
Harvard University
Cambridge, MA, USA
steven_finch@harvard.edu